

Chemical Analysis

Site:	Syntex - Vienna
ID #	100007452154
Break:	3.3
Other:	110

The available methods for determining 2,3,7,8-TCDD in ug/kg (ppb) or lower concentrations are ~~expensive~~, time-consuming, and difficult. The nature and use of these analyses necessitate elaborate quality control and quality assurance mechanisms (Clement and Stehl, 1973).

In the methods used, an isotopically labeled internal standard is added to the sample before the ~~extraction~~ step. Studies have shown that at 12 ppb or greater the final result is independent of the percent recovery of the internal standard. Other studies have shown that the optimum recovery of 2,3,7,8-TCDD from soil is ~~reached~~ only after several extractions. The internal standard may be recovered more efficiently because it is added in a solvent, and, consequently, it may not be bound as tightly to the soil. This latter combination would result in variable and oftentimes erroneously low results for the native 2,3,7,8-TCDD.

Another danger in adding recovery of an internal standard (which is added at 2.5 ppb) to be ~~is~~ ± 3., 35%, thereby recovering only the equivalent of 0.875 ppb of labeled 2,3,7,8-TCDD) is that the signal/noise ratio for a 1 ppb native 2,3,7,8-TCDD ~~will~~ would be decreased. If it is decreased to the extent that the signal representing 1 ppb of 2,3,7,8-TCDD (for 35% recovery from a 10-g sample, this ~~will~~ equal 3.5 ng in the final extract) is no longer within the linear range, ~~the~~ the percent recovery of the internal standard and the final result may ~~be~~ be independent. This is especially true for samples with a high background. Because of these reasons and the imprecision associated with other steps in the procedure, analytical results reflect a range rather than a specific number.



Recommendations for Further Study

A. Human

- Conduct additional case-control studies to determine the association between exposure to TCDD and related compounds and the incidence of soft tissue sarcomas.
- Conduct epidemiologic studies to characterize and follow populations exposed to TCDD. Determine their TCDD body burdens and the short- and long-term health effects.
- Include immunologic evaluation and evaluation of other likely target organs in epidemiologic studies of TCDD-exposed populations and develop baseline data for these end points.
- Develop better methods to analyze blood and tissue for TCDD at low concentrations. These should include radioimmunoassay (RIA) procedures and arylhydrocarbon hydroxylase (AHH) induction with biopsy material.
- Develop methods to increase excretion of TCDD after it accumulates in the body.
- Obtain better data on the toxicokinetics of TCDD in humans.
- Analyze human breast milk samples for TCDD and related compounds (to serve as an index for exposure and absorption) in exposed populations.

- Determine soil exposure/day by humans in various settings.

B. Experimental

- Conduct a series of studies related to the question of bioavailability of TCDD, to focus on differences by the type of soil, by routes of absorption, by species, by aging of soil/dioxin mixture, and by maturity of exposed animals.
- Study TCDD carcinogenicity and reproductive effects in the most sensitive animal species (guinea pig and/or subhuman primates). Carcinogenicity studies should also be conducted in non-sensitive and intermediate species to determine if the relation between the acute toxicity dose and the carcinogenicity dose is similar for all species.
- Conduct immunologic dose-response studies in young and adult animals.
- Conduct lifetime follow-up studies of immature exposed animals.
- Determine whether the ligand-receptor is integral to the mechanism(s) of action and search for antagonists and endogenous ligands in different tissues (including human cell systems in tissue culture).
- Determine whether TCDD exerts its carcinogenic potential by an initiator or promoter mechanism.
- Determine mechanisms for differences in species sensitivity.

- Study the mechanism of TCDD toxicity, particularly for cancer, as it relates to the ligand-receptor theory.
- Determine tissue TCDD levels in animals (guinea pigs) exposed *in situ* to TCDD-contaminated areas, thus providing further information on exposure and bioavailability.
- Prepare polychlorinated dibenzodioxin isomer standards for future laboratory work.
- Establish the role of metabolism more clearly.
- Establish the interactions of ligand-receptor complexes with macromolecules in hopes of ascertaining functional differences in these interactions between "sensitive" and "non-sensitive" species. These findings could be used in human risk assessment by providing a direct relationship between exposure and mechanism of toxicity.

C. Environmental

- Develop utilizable methods for detoxifying TCDD-contaminated soil.
- Determine ambient (background) levels of TCDD in soil, air, water, etc. (as well as in people).
- Establish "sentinel" animals as indicators of the bioavailability of TCDD in contaminated areas.

Table 1. Fetal loss in Rhesus Macaques after Oral Doses of TCDD during Weeks 4 through 6 of Pregnancy (McNulty, 1982)

Group	Total dose (ug/kg)	Schedule	Fetal losses	Gestational age of lost fetus (days)	Maternal toxicity
I	5	9 divided doses, 3 times a week	2/2	47, 50	2/2
	1	Same	3/4	50, 57, ?	1/4
	0.2	Same	1/4	?	0/4
II	1	1 dose, day 25	3/3	48, ?, ?	2/3
	1	1 dose, day 30	3/3	50, 51, 55	3/3
	1	1 dose, day 35	2/3	53, 108	1/3
	1	1 dose, day 40	2/3	100, 100	0/3
III	0	9 divided doses, 3 times a week	3/12	118, ?, ?	0/12

Table 2. Summary of Neoplastic Alterations observed in Sprague-Dawley Rats Fed Subacute Levels of TCDD for 78 Weeks (Van Miller et al., 1977)

Level of TCDD ng/kg	No. of animals with neoplasms*	No. of neoplasms	Diagnosis
0	0	0	--
1	0	0	--
5	5	6	1 ear duct carcinoma 1 lymphocytic leukemia 1 adenocarcinoma (kidney) 1 malignant histiocytoma (peritoneal)** 1 angiosarcoma (skin) 1 Leydig cell adenoma
50	3	3	1 fibrosarcoma (muscle) 1 squamous cell tumor (skin) 1 astrocytoma (brain)
500	4	4	1 fibrosarcoma (muscle) 1 carcinoma (skin) 1 adenocarcinoma (liver) 1 sclerosing seminoma (testes)
1000	4	5	1 cholangiocarcinoma (liver) 1 angiosarcoma (skin) 1 glioblastoma (brain) 2 malignant histocytomas (peritoneal)**
5000	7	10	4 squamous cell tumors (lung) 4 neoplastic nodules (liver) 2 cholangiocarcinomas (liver)

*10 animals per group.

**Metastases observed.

Table 3. Total and Individual Tumors in Treated Male and Female Rats of Significantly Different Incidence than Those in Nontreated Control Rats (Kociba et al., 1978)

<u>Tumor or tumor-like lesion</u>	<u>Dose level (ug TCDD/kg b.w./day)</u>									
	<u>0</u>		<u>0.001</u>		<u>0.01</u>		<u>0.1</u>			
	<u>M</u>	<u>F</u>	<u>M</u>	<u>F</u>	<u>M</u>	<u>F</u>	<u>M</u>	<u>F</u>	<u>M</u>	<u>F</u>
No. of rats examined	85	86	50	50	50	50	50	50	50	49
Hepatocellular neoplastic nodules	6	8	0	3	3	18 ¹	2	23 ¹		
Hepatocellular carcinoma	2	1	0	0	0	2	1	11 ¹		
Stratified squamous cell carcinoma of hard palate or nasal turbinates	0	0	0	0	0	1	7 ¹	4 ¹		
Keratinizing squamous cell carcinoma of lung	0	0	0	0	0	0	1	7 ¹		
Benign tumor of uterus	-	28	-	12	-	11	-	7 ²		
Benign neoplasm of mammary gland	-	73	-	35	-	36	-	24 ²		
Mammary carcinoma	-	8	-	4	-	4	0	0 ²		
Pituitary adenoma	26	43	6	18	11	13	13	12 ²		
Subcutaneous carcinomas	10	1	1 ²	1	5	0	6	0		
Acinar adenoma of pancreas	14	0	7	1	5	0	2 ²	1		
Adenoma of adrenal cortex	0	9	0	6	2	2	5 ¹	5		
Pheochromacytoma of adrenal gland	28	7	6	3	10	1	4 ¹	2		

Table 3. - continued

¹ Statistically greater than control data when analyzed by using the Fischer exact probability test, $p < 0.05$.

² Statistically less than control data when analyzed by using the Fischer exact probability test, $p < 0.05$.

Table 4. The Incidence of Dose-Related Tumors in Osborne-Mendel Rats and B6C3F1 Mice Treated with TCDD (NTP Study, 1982a and b).

Osborne-Mendel Rats			
Tumor Type	Dose ($\mu\text{g}/\text{kg}/\text{wk}$)	Sex	Incidence
Follicular-cell adenomas, thyroid	0	M	1/69
	0.01	M	5/48
	0.05	M	6/50
	0.5	M	10/50 ($P=0.001$)
Neoplastic nodules of the liver	0	F	3/73
	0.01	F	2/45
	0.05	F	1/49
	0.5	F	6/47
Neoplastic nodules or hepatocellular carcinoma	0	F	5/75
	0.01	F	1/49
	0.05	F	3/50
	0.5	F	12/49 ($P=0.006$)
B6C3F1 Mice	0	F	5/75
	0.01	F	1/49
	0.05	F	3/50
	0.5	F	14/49 ($P=0.001$)
B6C3F1 Mice			
Tumor type	Dose ($\mu\text{g}/\text{kg}/\text{wk}$)	Sex	Incidence
Follicular-cell adenomas of the thyroid	0	F	0/69
	0.04	F	3/50
	0.2	F	1/47
	2.0	F	5/46 ($P=0.009$)
Hepatocellular carcinoma	0	M	8/73
	0.01	M	9/49
	0.05	M	8/49
	0.5	M	17/50 ($P=0.002$)
Lymphoma and leukemia	0	F	1/73
	0.04	F	2/50
	0.2	F	2/48
	2.0	F	6/47 ($P=0.014$)

TABLE 5
Daily Soil Ingestion Patterns by Age

<u>Age Group</u>	<u>Soil Intested</u>
0-9 months	0 grams
9-18 months	1 gram
1 1/2-3 1/2 years	10 grams
3 1/2-5 years	1 gram
> 5 years	100 milligrams

TABLE 6

Daily Deposition of Soil on Skin by Age

<u>Age Group</u>	<u>Amount on Skin</u>
0-9 months	0 grams
9-18 months	1 gram
1 1/2-3 1/2 years	10 grams
3 1/2-15 years	1 gram
> 15 years	100 milligrams

Table 7: Important Lesion Sites in the Kociba TCDD Study

<u>Site:</u>	<u>Sex</u>	<u>Pathologist</u>	<u>Control</u>	<u>0.001</u>	<u>0.01</u>	<u>0.1</u>
Stratified squamous cell						
carcinoma of the tongue	Males	Kociba	0/76	1/49	1/49	3/42
		Squire	0/77	1/44	1/49	3/44
Nasal turbinates/hard						
palate squamous cell						
carcinoma	Males	Kociba	0/51	0/34	0/27	4/30
		Squire	0/55	1/34	0/26	6/30
	Females	Kociba	0/54	0/30	1/27	4/24
		Squire	0/54	0/30	1/27	5/22
Hepatoellular nodules						
and carcinoma	Females	Kociba	9/35	3/50	18/50	34/48
		Squire	16/35	8/50	27/50	33/47
Lung keratinizing						
squamous cell carcinoma	Females	Kociba	0/86	0/50	0/49	7/49
		Squire	0/86	0/50	0/49	8/47

Table 8: Estimates and Approximate 95% Lower Confidence Limits for the VSD[1] of TCDD from the Kociba Study in Sprague-Dawley Rats

<u>Lesion/Pathologist</u>	<u>Model[2]</u>	<u>Added Risk</u>		<u>Chi-squared</u> G-O-S[4]
		<u>1.E-4</u>	<u>1.E-6</u>	
♂				
Male Rat Stratified				
squamous cell carcinoma of				
the tongue	Kociba	Linear*	142581 (59430)	1426 (595) 1.43
	Squire	Linear*	151793 (62642)	1518 (626) 1.60
Male rat nasal turbinates				
or hard palate squamous				
cell carcinoma	Kociba	Cubic*	8877014 (17493)	1912466 (175) 0.00
		Linear	77507 (37566)	775 (376) 0.39
	Squire	Cubic*	7757813 (11576)	1671343 (117) 2.39
		Linear	49468 (26431)	495 (264) 2.72
Female rat lung keratinizing				
squamous cell carcinoma				
	Kociba	Cubic*	8659951 (32477)	1865707 (325) 0.01
		Linear	72600 (41318)	726 (413) 0.82
	Squire	Cubic*	8126063 (26834)	1750685 (268) 0.01
		Linear	60375 (35490)	604 (355) 0.98
Female rat nasal turbinates				
or hard palate squamous				
cell carcinoma	Kociba	Linear*	49771 (25812)	498 (258) 0.49
	Squire	Linear*	37237 (20313)	372 (203) 0.20

Table 8 - Continued

<u>Lesion/Pathologist</u>	<u>Model[2]</u>	<u>Added Risk</u>		<u>Chi-squared</u>
		<u>1.E-4</u>	<u>1.E-6</u>	<u>G-O-F[4]</u>
Female rat liver				
hepatocellular carcinoma or				
adenoma	Kociba	Linear*	7742 (5725)	77 (57.2) . 6.35
		Transf[3]	3836 (2919)	38 (29.1) 2.18
	Squire	Linear*	8649 (6074)	86 (60.7) 10.36
		Transf[3]	3760 (2763)	38 (27.6) 4.30

[1] Entries are virtually safe dose (VSD) Linear-cubic lower confidence bounds (LCB) in fg/kg b.w./day by using the multistage model.

[2] Linear Model: $P(d)=1-c(-a-b*d)$ Cubic Model: $P(d)=1-e(-a-c*d^{**3})$.

[3] A multistage model is fitted to the average liver concentration of TCDD, and safe doses are transformed by using the relationship:

$$\text{Administered dose} = \text{Liver dose}/510.297$$

The divisor is the least-squares slope between the administered and the average liver dose of TCDD for control and the two lowest doses.

[4] G-O-F = goodness of fit.

*best fit.

Table 9: Estimates and Approximate 95% Lower Confidence Limits for the VSD[1] of TCDD from the NCI/NTP Study

Lesions:	Added Risk		Chi-squared	
	Model[2]	1.E-4	1.E-6	G-O-F [3]
Male rat thyroid				
Follicular-cell adenoma	Linear*	40291 (21435)	403 (214)	4.81
Female rat thyroid				
Follicular-cell adenoma	Cubic*	7142868 (36711)	1542868 (367)	0.48
	Linear	75737 (34895)	757 (349)	0.79
Female rat liver				
Neoplastic nodules	Linear*	1442863 (19298)	105571 (193)	1.31
	Linear	31520 (18764)	315 (188)	1.61
Nodules and Carcinomas	Quadratic*	1338200 (16523)	133820 (165)	1.31
	Linear	25552 (16001)	256 (160)	1.74
Male Mouse Liver				
Hepatocellular Carc.	Linear*	26293 (15147)	263 (151)	1.01
Adenomas and Carc.	Linear*	13394 (8633)	134 (86)	0.14
Female Mouse Liver				
Hepatocellular Carc.	Linear*	246021 (121594)	2460 (1216)	0.74
Adenomas and Carc.	Linear*	145013 (77376)	1450 (779)	2.59
Female Mouse Thyroid				
Follicular-Cell Adenoma	Linear*	301051 (139090)	3011 (1391)	3.75

Table 9 - Continued

<u>Lesions:</u>	<u>Model[2]</u>	<u>Added Risk</u>		<u>Chi-squared</u> <u>G-O-F [3]</u>
		<u>1.E-4</u>	<u>1.E-6</u>	
Female Mouse Sarcoma				
Subcutaneous Tissue	Lin-Cubic*	430263 (142303)	4303 (1428)	0.05
	Linear	295289 (142777)	2953 (1427)	0.06
Female Mouse Lymphoma and Leukemia	Linear*	102852 (54271)	1028 (543)	0.02

[1] Entries are virtually safe dose (VSD) lower confidence bounds (LCB) in fg/kg b.w./day by using the multistage model.

[2] Linear: $P(d)=1-3(-a-b*d)$ Quad: $P(d)=1-3(-a-c*d**2)$

Cubic: $P(d)=1-3(-a-f*d**3)$ Lin-Quadratic: $P(d)=1-3(-a-b*d-c*d**2)$

Lin-Cubic: $P(d)=1-3(-a-b*d-f*d**3)$

[3] G-O-F = goodness of fit.

* best fit.

Table 10: Concentrations of TCDD in Soil That Are Projected to Produce the Maximum Allowable Residues in Foods

Food	TCDD in fat, pg/g	Observed Ratio ¹	Soil, pg/g
Beef ²	7.9	0.39	20
Beef(cull dairy) ³	7.9	0.10	79
Pork	22.7	1.86	12
Milk	2.5	0.40	6.2

¹ Concentration of polybrominated biphenyls (PBB) in product/concentration of PBB in soil (Fries and Jacobs, 1983).

² Includes dairy cattle that have never lactated.


³ Older cows. Younger cows would approach the values for beef cattle.

Table 11: Daily Consumption, Fat Content and Allowable Concentration of TCDD in Fat to Maintain Human TCDD Intake Below 100 pg/day/person

	Consumption, g/day	Fat, %	Fat consump- tion, g/day	Allowable TCDD in fat, pg/g¹
Beef	105	8-12	8.4-12.6	7.9
Pork	54	6-8	3.2-4.3	22.7
Milk	1000	4	40	2.5

¹ Based on the higher fat intake value.

ESTIMATED AVERAGE DAILY DOSE CORRESPONDING TO INITIAL TCDD - SOIL CONTAMINATION LEVELS



EXCESS LIFETIME RISK OF DEVELOPING CANCER CORRESPONDING TO INITIAL TCDD - SOIL CONTAMINATION LEVELS

FIGURE 2





FIGURE 3



Legends for Figures

- Figure 1** This figure illustrates the average daily dose of TCDD which would be received if 100, 10, or 1% of the accessible soil were contaminated with the peak recorded level of TCDD. The boundaries for an excess lifetime cancer risk for 10^{-5} and 10^{-6} are also illustrated.
- Figure 2** This figure illustrates the lower and upper bounds of the virtually safe dose for a lifetime excess cancer risk. This information was derived from the calculations developed from animal data (Table 8 and 9). At the concentration of 1 ppb if 100% of the accessible soil contained TCDD at this concentration the area of risk which is bordered by the upper and lower bound of the virtually safe dose does not represent an unacceptable cancer risk given the fact that the background cancer incidence in the general population is of a much higher order of magnitude. If less than 100% of the soil was contaminated this risk would even be further reduced. However, at levels much above 1 ppb the risk would become unacceptable.
- Figure 3** Surface area covered by 1 and 10 g of dirt less than 1 mm thick.
- Figure 4** Average surface area of the palm of one hand of 2- and 4-year old children, adult females and males.

Appendix I

CONSULTANTS

Conference on Polyhalogenated Aromatic Compounds

Atlanta, GA.

June 28-29, 1983

Dr. Donald Barnes
Senior Science Advisor for Pesticides
and Toxic Substances
Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460

Dr. Judy Bellin
Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460

Dr. Frank Cordle
Food and Drug Administration
HFF-103
200 C Street, SW
Washington, D.C. 20204

Dr. James Falco
Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460

Dr. George F. Fries
Research Animal Scientist
Pesticide Degradation Lab
U.S. Department of Agriculture
Beltsville, Maryland 20705

Dr. Donald Grant
Health Protection Branch
Health and Welfare Canada
Health Protection Branch Building
Tunney Pasture
Ottawa, Ontario K1A0L2

Dr. Robert Harris
Co-Director of Hazardous Waste
and Research Division
Center for Energy and Environmental Studies
Princeton University
Von Neumann Building
Princeton, New Jersey 08544

Dr. David G. Noel
National Institute of Environmental Health Sciences
P.O. Box 12233
Research Triangle Park, North Carolina 27709

Dr. Nancy Kim
Director of Bureau of Toxic Substances Assessment
New York State Department of Health
Empire State Plaza
Tower Building, 14th Floor
Albany, New York 12237

Dr. George Lucier
National Institute of Environmental Health Sciences
P.O. Box 12233
Research Triangle Park, North Carolina 27709

Dr. Norton Nelson
Professor of Environmental Medicine
New York University Medical Center
Institute of Environmental Medicine
500 First Avenue
New York, New York 10016

Dr. Henry Pitot
Director, McArdle Cancer Research Institute
University of Wisconsin
Madison, Wisconsin 53706

Dr. Charles F. Reinhardt
DuPont Company
Haskell Lab
P.O. Box 850, Elkton Road
Newark, Delaware 19711

Dr. Robert G. Tardiff
1423 Trapline Court
Vienna, Virginia 22180

Dr. John Van Ryzin
Professor, Division of Biostatistics
School of Public Health
Columbia University
600 West 118 Street
New York, New York 10032

CENTERS FOR DISEASE CONTROL

Dr. Vernon N. Houk
Director,
Center for Environmental Health

Dr. David D. Bayse
Director,
Clinical Chemistry Division
Center for Environmental Health

Dr. Henry Falk
Chief, Special Studies Branch
Chronic Diseases Division
Center for Environmental Health

Dr. Renate D. Kimbrough
Medical Officer
Center for Environmental Health

Dr. Larry K. Lowry
Centers for Disease Control
National Institute for Occupational Safety and Health
Robert Taft Laboratory
4676 Columbia Parkway
C-26
Cincinnati, Ohio 45226

Ms. Elizabeth Ward
Epidemiologist (replacing Dr. Fingerhut)
Centers for Disease Control
National Institute for Occupational Safety and Health
Robert Taft Laboratory
4676 Columbia Parkway
F-6
Cincinnati, Ohio 45226

Dr. Paul J. Wiesner
Director,
Chronic Diseases Division
Center for Environmental Health

Appendix II

Date: August 1, 1983
 From: Mathematical Statistician, BRAP
 Subject: Teratogenic and Reproductive Effects of 2,3,7,8-TCDD as Noted by Murray et al.
 To: Director, BRAP

I have previously reviewed the article by Murray et al. entitled "Three generation reproduction study of rats given 2,3,7,8-Tetrachlorodibenzo-p-dioxin in the diet." I must report that the analysis of the results from this study are suspect. Several reasons lead me to this conclusion. First, in a reanalysis of these data using trend tests instead of paired comparisons, the results were different (Nisbet and Paxton, Am. Stat., 36[3]. Most notably, Nisbet and Paxton concluded that the 0.001 dose level (not the 0.01 dose level as is stated in the synopsis from Kimbrough) is not a NOAEL, whereas Murray et al. concluded 0.001 was a NOAEL. In addition, in the table below, you will note that the fertility index increases with each subsequent generation in the control and low dose groups, but remains constant in the 0.01 dose group. Murray et al. and Nisbet and Paxton both conclude there is decreased fertility in the 0.01 dose group as compared to the control group. Without more information concerning what is a normal fertility index in these animals and just based upon these data, I cannot agree with their result. Including these remarks and noting the difficulty of the design for analysis, I would conclude there is insufficient evidence for an effect at the 0.01 dose.

C.J. Portier

TABLE I
Fertility Index (FI[1] of Rats Given TCDD

Generation	D	Micrograms TCDD/kg/day							
		0		0.001		0.01		0.1	
		FI	%	FI	%	FI	%	FI	%
f0	first mating	14/32	44	10/20	50	12/20	60	3/31	10
	second mating	21/32	66	14/19	74	15/20	75	1/30	3
f1		22/26	85	15/17	88	12/21	57		
f2		28/32	88	20/20	100	11/20	55		

[1] The fertility index used in this table is the number of females delivering a litter divided by the number of females placed with a male.

References

Allen, J.R., Barsotti, D.A., Lambrecht, L.K., and Van Miller, J.P. 1979.

Reproductive effects of halogenated aromatic hydrocarbons on non-human primates. Ann. N.Y. Acad. Sci. 320:419-425.

Ames, B.N. 1983. Dietary carcinogens and anticarcinogens oxygen radicals and degenerative diseases. Science 222:1256-1264.

Anonymous. 1983. EPA, Dow clash on dioxin in Michigan River. Environ. Health Lett. (Fishbein, G.W., ed.) April 1:8.

Appelgren, L.E., Brandt, T., Britteba, E.B., Gilmer, M., and Gustafsson, J.A. 1983. Autoradiography of 2,3,7,8-tetrachloro¹⁴cl-dibenzo-p-dioxin (TCDD): Accumulation in the nasal mucosa. Chemosphere 12:545-548.

Bauer, H., Schulz, K.H., and Spiegelberg, U. 1961. Berufliche Vergiftungen bei der Herstellung von Chlorphenol-Verbindungen. Arch. Gewerbepathol. Gewerbehyg. 18:538-555.

Bleiberg, J., Wallen, M., Brodkin, R., and Applebaum, I.L. 1964. Industrially acquired porphyria. Arch. Dermatol. 89:793-797.

Bumb, R.R., Crummett, W.B., Cutie, S.S., Gledhill, J.R., Hummel, R.H., Kage, R.O., Lamparski, L.L., Luoma, E.V., Miller, D.L., Nestrick, T.J., Shadoff, L.A., Stehl, R.H., and Woods, J.S. 1980. Trace chemistries of fire: A source of chlorinated dioxins. Science 210:385-390.

Buser, H.R., Bosshardt, H.P., and Rappe, C. 1973. Identification of polychlorinated dibenzo-p-dioxin isomers found in fly ash. *Chemosphere* 7:165-172.

Carlstedt-Duke, J., Elfstrom, G., Snochowski, M., Hogberg, B., and Gustafsson, J.A. 1978. Detection of the 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) receptor in rat liver by isoelectric focusing in polyacrylamide gels. *Toxicol. Lett.* 2:365-373.

Carter, C.D., Kimbrough, R.D., Liddle, J.A., Cline, R.F., Zack, M.M. Jr., Barthel, W.F., Koehler, R.E., and Phillips, P.E. 1975. Tetrachlorodibenzodioxin: An accidental poisoning episode in horse arenas. *Science* 138:738-40.

Coggon, D., Acheson, E.D. 1982. Do phenoxy herbicides cause cancer in man? *Lancet* 7:1057-1059.

Consultative Council on Congenital Abnormalities in the Yarram District. 1978. Ministry of Health, Melbourne, Australia.

Cook, R.R. 1981. Dioxin, chloracne, and soft tissue sarcoma. *Lancet* 618:3-14.

Cook, R.R., Townsend, J.C., Ott, M.G., and Silverstein, L.G. 1980. Mortality experience of employees exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *J. Occup. Med.* 22(8):530-532.

Crummett, W.B., and Stchl, R.H. 1973. Determination of chlorinated dibenzo-p-dioxins and dibenzofurans in various materials. Environ. Health Perspect. 5:15-25.

Crump, K., Guess, H.A., and Deal, K.I. 1977. Confidence intervals and test of hypothesis concerning dose response relations inferred from animal carcinogenicity data. Biometrics 33:437-51.

DiDomenico, A., Silano, V., Viviano, G., and Zapponi, G. 1980. Accidental release of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) at Seveso, Italy. Ecotoxicol. Environ. Safety 4:346-356.

Eiceman, G.A., Viau, A.C., and Karasek, F.W. 1980. Ultrasonic extraction of polychlorinated dibenzo-p-dioxins and other organic compounds from fly ash from municipal incinerators. Anal. Chem. 52:1492-1496.

Eriksson, M., Hardell, L., Berg, N., Moller, T., and Axelson, O. 1981. Soft tissue sarcomas and exposure to chemical substances: A case referent study. Br. J. Ind. Med. 38:27-33.

Faith, R.E., and Luster, M.I. 1979. Investigations on the effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) on parameters of various immune functions. Ann. N.Y. Acad. Sci. 320:564-571.

Faith, R.E., Luster, M.I., and Moore, J.A. 1978. Chemical separation of helper cell function and delayed hypersensitivity responses. *Cell. Immunol.* 40:275-284.

Farber, E. 1974. Pathogenesis of liver cancer. *Arch. Pathol.* 98:145-148.

Fillipini, G., Bordo, B., Crenna, P., Massetto, N., Musicco, M., and Boeri, R. 1981. Relationship between clinical and electrophysiological findings and indicators of heavy exposure to 2,3,7,8-tetrachloro-dibenzodioxin. *Scand. J. Work Environ. Health.* 7:257-62.

Fingerhut, M.A., Halperin, W.E., Houchar, P.A., Smith, A.B., Groth, D.H., and Russell, W.O. 1984. Review of Exposure and Pathology Data for Seven Cases Reported Among Soft Tissue Sarcoma Among Persons Occupationally Exposed to Dioxin-Contaminated Herbicides. (Paper in preparation.)

Firestone, D., Clower, M., Borsetti, A.P., Teske, R.H., and Long, P.E. 1979. Polychlorodibeno-p-dioxin and pentachlorophenol residues in milk and blood of cows fed technical pentachlorophenol. *J. Agric. Food Chem.* 27:1171.

Fries, G.F. 1982. Potential polychlorinated biphenyl residues in animal products from application of contaminated sewage sludge to land. *J. Environ. Qual.* 11:14.

Fries, G.F., and Jacobs, L.W. 1983. Tissue residues in livestock confined to unpaved lots with soil that contained polybrominated biphenyl. *J. Anim. Sci.* (Abstract in press).

Fries, G.F., and Marrow, G.S. 1975. Excretion of polybrominated biphenyls in milk of cows. J. Dairy Sci. 58:947.

Fries, G.F., and Marrow, G.S. 1982. Soil ingestion by swine as a route of contaminant exposure. Environ. Toxicol. 1:201-204.

Casiewicz, T.A., and Neal, R.A. 1979. 2,3,7,8-tetrachlorodibenzo-p-dioxin tissue distribution excretion and effects on clinical chemical parameters in guinea pigs. Toxicol. Appl. Pharmacol. 5:329-340.

Goldstein, J.A. 1983. Structure activity relationships for the biochemical effects and the relationship to toxicity (Chapter 6). In Halogenated Biphenyls, Terphenyls, Napthalenes, Dibenzodioxins and Related Products, ed. R.D. Kimbrough, pp. 151-190. Topics Environ. Health 4. New York: Elsevier, North Holland Biomedical - Amsterdam.

Greenlee, W.F., and Poland, A. 1979. Nuclear uptake of 2,3,7,8-tetrachloro-dibenzo-p-dioxin in C57BL/6J and DBA/2J mice. J. Biol. Chem. 254:9814-9821.

Hardell, L., Eriksson, M., Lennar, P., and Lundgren, E. 1981. Malignant lymphoma and exposure to chemicals especially organic solvents chlorophenols and phenoxy acids: A case-control study. Br. J. Cancer 43(2):169-176.

Hardell, L., and Sandstrom, A. 1979. Case-control study: Soft-tissue sarcomas and exposure to phenoxyacetic acids and chlorophenols. Br. J. Cancer 39:711-717.

Hay, A. 1976. Toxic cloud over Seveso. Nature 262:636-3.

Hay, A. 1982. The Chemical Scythe. New York and London: Plenum Press.

Honchar, P.A., and Halperin, W.E. 1981. 2,4,5-trichlorophenol and soft tissue sarcoma. Lancet. 1(8214):268-269.

International Agency for Research on Cancer (IARC). 1977. Chlorinated dibenzodioxins. IARC Monog. Eval. Carcino. Risk Mon 15:41-102.

Jensen, N.E., and Walker, A.E. 1972. Chloracne: Three cases. Proc. R. Soc. Med. 65:687-688.

Jirasek, L., Kalensky, J., Kubec, K., Pazderova, J., and Lukas, E. 1976. Chloracne, Porphyria Cutanea Tarda und andere Intoxikationen durch Herbizide. Hautarzt 27:328-33.

Jones, C.; and Butler, W.H. 1974. A morphological study of the liver lesion induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin in rats. J. Pathol. 112:93-97.

Khera, K.S., and Suddick, J.A. 1973. Polychlorodibenzo-p-dioxins: Perinatal effects and the dominant lethal test in Wistar rats. Adv. Chem. Ser. 120:70-84.

Kimbrough, R.D. (ed.). 1980. Halogenated biphenyls, terphenyls, naphthalenes, dibenzodioxins and related products. Top. Environ. Health 4:406.

Kimbrough, R.D., Carter, C.D., Liddle, J.A., Cline, R.E., and Phillips, P.E. 1977. Epidemiology and pathology of a tetrachlorodibenzodioxin poisoning episode. Arch. Environ. Health 32:77-85.

Kitchin, K.T., and Woods, J.S. 1979. 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) effects on hepatic microsomal cytochrome p-448 mediated enzyme activities. Toxicol. Appl. Pharmacol. 47:537-546.

Kociba, R.J., Keyes, D.G., Beyer, J.E., Carreson, R.M., Wade, E.E., Dittenber, D.A., Malmins, R.P., Frauson, L.F., Park, D.N., Barnard, S.D., Hummel, R.A., and Humiston, C.G. 1978. Results of a two year chronic toxicity and oncogenicity study of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in rats. Toxicol. Appl. Pharmacol. 46:279-303.

Kociba, R.J., and Schwetz. 1982. Toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). Drug Metabol. Reviews. 13:387-406.

Kooke, R., Lustenhouwer, J.W.A., Olie K., and Hutzinger, O. 1981. Extraction efficiencies of polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans from fly ash. Anal. Chem. 53:461-463.

Krasovisky, G.N. 1976. Extrapolation of experimental data from animals to man. Environ. Health. Perspect. 13:51-58.

Manara, L., Coccia, P., Croci, T. 1982. Persistent tissue levels of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in the mouse and their reduction as related to prevention of toxicity. *Drug. Metab. Rev.* 13:423-446.

Manis, J., and Kim, G. 1980. Polybrominated biphenyl: Acute and chronic effect on iron absorption and benzpyrene hydroxylase. *Toxicol. and Appl. Pharmacol.* 54:41-47.

May, G. 1973. Chloracne from the accidental production of tetrachlorodibenzodioxin. *Br. J. Ind. Med.* 30:276-283.

May, G. 1982. Tetrachlorodibenzodioxin: A survey of subjects ten years after exposure. *Br. J. Ind. Med.* 39:128-135.

McConnell, E.E., Moore, J.A., and Dalgard, D.W. 1978. Toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in rhesus monkeys (*Macaca mulatta*) following a single oral dose. *Toxicol. Appl. Pharmacol.* 43:175-187.

McConnell, E.E., Moore, J.A., Haseman, J.K., and Harris, M.W. 1978. The comparative toxicity of chlorinated dibenzo-p-dioxins in mice and guinea pigs. *Toxicol. Appl. Pharmacol.* 44:335-56.

McConnell, E.E., et al. 1983. Studies on the bioavailability of dioxin in soil. NIEHS, Research Triangle Park, N.C. (personal written communication)

McNulty, W.P. 1982. Rhesus macaques: Pertinence for studies on the toxicity of chlorinated hydrocarbon environmental pollutants. In Advanced Views in Primate Biology, eds. A.B. Chiarelli and R.S. Carruccini, pp. 111-113. Berlin: Springer-Verlag.

Milby, T.H., Husting, E.L., Whorton, M.D., and Larson, S. 1980. Potential health effects associated with the use of phenoxy herbicides: A summary of recent scientific literature: A report for the National Forest Products Association. Berkeley: Environmental Health Associates.

Milham, S., Jr. 1982. Herbicides, occupation, and cancer. Lancet 1:1464-1465.

Moore, J.A., Harris, M.W., and Albro, P.W. 1976. Tissue distribution of (¹⁴C) tetrachlorodibenzo-p-dioxin in pregnant and neonatal rats. Toxicol. Appl. Pharmacol. 37:146-147.

Murray, F.J., Smith, F.A., Nitschke, K.D., Humiston, C.G., Kociba, R.J., and Schwetz, B.A. 1979. Three generation reproduction study of rats given 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in the diet. Toxicol. Appl. Pharmacol. 50:241-252.

National Research Council of Canada (NRCC). 1981. Polychlorinated dibenzo-p-dioxins. Publ. NRCC No. 18574 of the Environmental Secretariat. Publ. Ottawa, Canada: NRCC.

National Toxicology Program (NTP). 1982a. Carcinogenesis bioassay of 2,3,7,8-tetrachlorodibenzo-p-dioxin (CAS Nol 1746-01-6) in Swiss-Webster mice (dermal study). NTP Tech. Rep. Ser. Issue 201:113. NTP.

National Toxicology Program (NTP). 1982b. Carcinogenesis bioassay of 2,3,7,8-tetrachlorodibenzo-p-dioxin (CAS Nol 1746-01-6) in Osborne-Mendel rats and B6C3F1 mice (gavage study). NTP Tech. Rep. Ser. Issue 209:195. NTP.

Nau, H., and Bass, R. 1981. Transfer of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) to the mouse embryo and fetus. Toxicol. 20:299-308.

Neal, R.A., Olson, J.R., Gasiewicz, T.A., and Geiger, L.E. 1982. The toxicokinetics of 2,3,7,8-tetrachlorodibenzo-p-dioxin in mammalian systems. Drug Metabbl. Rev. 13:355-385.

Nesbit, I.C.T, and Paxton, M.B. 1982. Statistical aspects of 3 generation studies of the reproductive toxicity of TCDD and 2,4,5-T. Am. Stat. 36:290-298.

Morstrom, R.J., Hallett, D.J., Simon, M., and Mulvihill, M.J. 1982. Analysis of Great Lakes herring gull eggs for tetrachlorodibenzo-p-dioxins. In Chlorinated Dioxins and Related Compounds: Impact on the Environment, eds. O. Hutzinger, R. W. Frei, E. Merian, and F. Pocchiari, Vol. 5, pp. 173-181. New York: Pergamon.

Office on Smoking and Health. 1982. The health consequences of smoking: Cancer. A report of the Surgeon General. Rockville, Maryland: Public Health Service, United States Department of Health and Human Services.

Oliver, R.M. 1975. Toxic effects of 2,3,7,8 tetrachlorodibenzo 1,4 dioxin in laboratory workers. Br. J. Ind. Med. 32:49-53.

Olson, J.R., Gasiewicz, T.A., and Neal, R.A. 1980. Tissue distribution, excretion and metabolism of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in the golden syrian hamster. Toxicol. Appl. Pharmacol. 56:78-85.

Ott, M.G., Holder, B.B., and Olson, R.D. 1980. A mortality analysis of employees engaged in the manufacture of 2,4,5-trichlorophenoxyacetic acid. J. Occup. Med. 22(1):47-50.

Pazderova-Vejlupkova, J., Lucas, E., Nemcova, M., Pickova, J., and Jirasek, L. 1981. The development and prognosis of chronic intoxication by tetrachlorodibenzo-p-dioxin in men. Arch. Environ. Health 36:5-11.

Pinkerton, H., in Anderson, W.A.D., and Kissaw, J.M. 1977. Pathology. pp. 625-628. Saint Louis: CV Mosby Co. .

Piper, W.N., Rose, R.Q., and Gehring, P.J. 1973. Excretion and tissue distribution of 2,3,7,8-tetrachlorodibenzo-p-dioxin in the rat. Environ. Health Perspect. 5:241-244.

Pitot, H.C., Goldworthy, T., Campbell, H.A., and Poland, A. 1980. Promotion by 2,3,7,8-tetrachlorodibenzo-p-dioxin of hepatocarcinogenesis from diethylnitrosamine. *Cancer Res.* 40:3616-3620.

Pocchiani, F., Silano V., and Zampieri, A. 1979. Human health effects from accidental release of TCDD at Seveso (Italy)., *Ann. NY. Acad. Sci.* 320:311.

Poiger, H., Busser, H.R., Weber, H., Zweifel, U., and Schlatter, C. 1982. Structure elucidation of mammalian TCDD metabolites. *Experientia* 38:434-486.

Poiger, H., and Schlatter, C. 1980. Influence of solvents and absorbance on dermal and intestinal absorption of TCDD. *Ford Cosmet. Toxicol.* 18:477-481.

Poland, A., and Glover, E. 1973. Chlorinated dibenzo-p-dioxins: Potent inducers of S-aminolevulinic acid synthetase and aryl hydrocarbon hydroxylase. II. A study of the structure activity relationship. *Mol. Pharmacol.* 9:736-747.

Poland, A., and Glover, E. 1975. Genetic expression of aryl hydrocarbon hydroxylase by 2,3,7,8-tetrachlorodibenzo-p-dioxin: Evidence for a receptor mutation in genetically nonresponsive mice. *Mol. Pharmacol.* 11:389-398.

Poland, A., and Glover, E. 1979. An estimate of the maximum in vitro covalent binding of 2,3,7,8-tetrachlorodibenzo-p-dioxin to rat liver protein ribosomal RNA and DNA. *Cancer Res.* 39:3341-3344.

Poland, A., Glover, E., and Kende, A.S. 1976. Stereoselective high affinity binding of 2,3,7,8-tetrachlorodibenzo-p-dioxin by hepatic cytosol. J. Biol. Chem. 251:4926-4946.

Poland, A., Palen, D., and Glover, E. 1982. Tumor promotion by TCDD in skin of BKS/hairless mice. Nature 300:271-273.

Poland, A.P., Smith, D., Matter, G., and Possick, P. 1971. A health survey of workers in a 2,4-D and 2,4,5-T plant. Arch. Environ. Health 22:316-327.

Ramsey, J.C., Hefner, J.G., Karbowski, R.J., Braun, W.H., and Gehring, P.J. 1979. The in vivo biotransformation of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in the rat (Abstract). Toxicol. Appl. Pharmacol. 48:A162.

Reggiani, G. 1980. Acute human exposure to TCDD in Seveso, Italy. J. Toxicol. Environ. Health 6:27-43.

Rogers, A.M., Anderson, M.E., and Back, K.C. 1982. The toxicity and mutagenicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin and similar acting compounds. Environ. Mutagenesis 4:391.

Q

Rose, J.Q., Ramsey, J.C., Wentzler, T.H., Hummel, R.A., and Gehring, P.J. 1976. The fate of 2,3,7,8-tetrachlorodibenzo-p-dioxin following single and repeated oral doses to the rat. Toxicol. Appl. Pharmacol. 36(2):209-226.

Smith, A.H., Fisher D.O., Gibbs, H.J., and Pearcey, H. 1982. The New Zealand soft tissue sarcoma case-control study: Interview findings concerning polychlorinated acid exposure. *Chemosphere* 12:565-571.

Smith, F.A., Schwetz, B.A., and Mitschke, K.D. 1976. Teratogenicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in CP-I mice. *Toxicol. Appl. Pharmacol.* 33:517-523.

Stohs, S.J., Hassan, H.Q., and Murray, W.J. 1983. Lipid peroxidation as a possible cause of TCDD toxicity. *Biochem. Biophys. Res. Commun.* 111:854-859.

Sweeney, G.D., and Jones, K.G. 1983. Studies of the mechanism of action of hepatotoxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) and related compounds. *Environ. Sci. Res.* 26:415-422.

Sweeney, G.D., Jones, K.G., Cole, F.M., Basford, D., and Krestynski, P. 1979. Iron deficiency prevents liver toxicity of 2,3,7,8 TCDD. *Science* 204:332-335.

Thibodeau, M., Freret-Lagye, R., and Link, R. 1972. Mortality study of persons exposed to dioxin in a trichlorophenoxy process accident that occurred in the LASF AG on Nov. 17, 1953. *Am. J. Ind. Med.* 3:179-189.

Thigpen, J.E., Faith, R.E., McConnell, E.E., and Moore, J.A. 1975. Increased susceptibility to bacterial infection as a sequela of exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Infect. Immunol.* 12:1319-1324.

Thomas, P.T., and Hinsdill, R.D. 1979. The effect of perinatal exposure to tetrachlorodibenzo-p-dioxin on the immune response in mice. *Drug Chem. Toxicol.* 2:77-98.

Toth, K., Somfai-Kelle, S., Sugar, J., and Bruce, J. 1979. Carcinogenicity testing of the herbicide 2,4,5-trichlorophenoxy ethanol containing dioxin and of pure dioxin in Swiss mice. *Nature* 278:548-549.

Van Miller, J.P., Lalich, J.J., and Allen, J.R. 1977. Increased incidence of neoplasms in rats exposed to low levels of 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Chemosphere* 10:625-32.

Vecchi, A., Montavalli, A., Sironi, M., Luini, W., Cairo, M., and Garattini, S. 1980. Effect of acute exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin on humoral antibody production in mice. *Chem. Biol. Interact.* 30:337-42.

Vos, J.G., Kreenitberg, J.G., Engel, H.W., Minderhoud, A., and van Voorle-Jansen, L.M. 1973. Studies on 2,3,7,8-TCDD-induced immune suppression and decreased resistance to concentrations and effect of thymosin treatment. *Toxicology* 5:75-86.

Vos, J.G., and Moore, J.A. 1974. Suppression of cellular immunity in rats and mice by maternal treatment with 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Int. Arch. Allergy Appl. Immunol.* 47:777-94.

Vos, J.G., Moore, J.A., and Zinkl, J.G. 1973. Effect of
2,3,7,8-tetrachlorodibenzo-p-dioxin on the immune system of laboratory
animals. Environ. Health Perspect. 5:149-62.

D

Zack, J.A., and Caffey, W.R. 1983. A mortality study of workers employed at
the Monsanto Company plant in Nitro, West Virginia. Environ. Sci. Res.
26:575-591.

Zack, J.A., and Suckind, R.R. 1980. The mortality experience of workers
exposed to tetrachlorodibenzodioxin in a trichlorophenol process accident. J.
Occup. Med. 22:11-4.

Selected References:

Bailey, D.L., and Tinkle, E.S. 1979. Particulate deposition from dry unpaved roadways. *Atmospheric Environment*. Great Britain: Pergamon. 13:661-668.

Cannoni, I., DiMuccio, A., Pontecorvo, D., Taggi, F., and Vergori, L. 1982. Laboratory investigation for the microbiological degradation of 2,3,7,8-tetrachlorodibenzo-p-dioxin in soil by addition of organic compost. In *Chlorinated Dioxins and Related Compounds: Impact on the Environment*, eds. O. Hutzinger, R.W. Frei, E. Merian, and F. Pocchiari, Vol. 5, pp. 95-103. New York:Pergamon.

Crow, K.D. 1982. Chloracne. *Seminars in Dermatology*. 1(4):305-314.

DiDomenico, A., Viviano, G., and Zapponi, G. 1982. Environmental persistence of 2,3,7,8-TCDD at Seveso. In *Chlorinated Dioxins and Related Compounds Impact on the Environment*, eds. O. Hutzinger, R.W. Frei, E. Merian, and F. Pocchiari, Vol. 5, pp. 105-114. New York:Pergamon.

Fanelli, R., Chiabrando, C., and Bonaccorsi, A. 1982. TCDD contamination in the Seveso incident. *Drug Metabolism Reviews* 13(3):407-422.

Harless, R.L., Oswald, E.O., and Lewis, R.G. 1982. Determination of 2,3,7,8,-tetrachlorodibenzo-p-dioxin in fresh water fish. *Chemosphere* 11(2):193-198.

Hutter, R., and Philippi, M. 1982. Studies on microbial metabolism of TCDD under laboratory conditions. In *Chlorinated Dioxins and Related Compounds: Impact on the Environment*, eds. O. Hutzinger, R.W. Frei, E. Merian, and F. Pocchiari, Vol. 5, pp. 87-97. New York:Pergamon.

Kimble, B.J., and Gross, H.L. 1980. Tetrachlorodibenzo-p-dioxin quantitation in stack-collected coal fly ash. *Science* 207:59-61.

Kociba, R.J., and Schwetz, B.A. Toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). *Drug Metabolism Reviews* 13(3):387-406.

Lee, I.P., Suzuki, K., and Nagayama, J. 1981. Metabolism of benzo (a) pyrene in rat prostate glands following 2,3,7,8-tetrachlorodibenzo-p-dioxin exposure. *Carcinogenesis* 2(9):823-31.

McMulty, W.P., and Nielsen, K.A. 1982. Persistence of TCDD in monkey adipose tissue. *Chem. Toxicol.* 20:985-987.

McMulty, W.P., Pomerantz, I.M., and Farrall, T.J. 1982. Chronic toxicity of 2,3,7,8-tetrachlorodibenzofurans for rhesus macaques. In Chlorinated Dioxins and Related Compounds Impact on the Environment, eds. O. Mutschler, B.W. Frei, E. Marais, and F. Pecchiali, Vol. 5, pp. 437-47. New York:Pergamon.

Poland, A., and Kende, A. 1975. 2,3,7,8-tetrachlorodibenzo-p-dioxin: Environmental contaminant and molecular probe. *Fed. Proc.* 35(12):2404-2411.

Van Looyen, M.J., Gupta, B.N., McConnell, E.E., and Moore, J.A. 1981. The influence of malnutrition on the toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in rats. *Toxicol.* 21:77-78.

Vejlupkona, J.P., Likus, E., Nemcova, M., Pickona, J., and Jirasek L. 1981.
The development and prognosis of chronic intoxication by tetrachloro-
dibenzo-p-dioxin in man. Arch. Environ. Health 36(1):5-11.

Vincent, J.H., and Abramruster, L. 1981. On the quantitative definition of
the inhalability of airborne dust. Ann. Occup. Hyg. 24(2):245-243.

Wipf, H.K., Homberger, E., Neuner, N., Ranalder, U.B., Vetter, W., and
Vuilleumier, J.P. 1982. TCDD-levels in soil and plant samples from the
Seveso area. In Chlorinated Dioxins and Related Compounds: Impact on the
Environment, eds. O. Nutzinger, and R.W. Frei, E. Merian, and F. Pocchiari,
Vol. 5, pp. 115-126. New York:Pergamon.

Acknowledgment

The statistical analysis of the rodent carcinogenicity data was performed by Dr. Christopher Portier, Biometry and Risk Assessment Program, National Institute of Environmental Health Sciences, P.O. Box 12233, Research Triangle Park, North Carolina 27709.

Dr. Paul J. Wiesner, David L. Forney, Linda W. Anderson, and Debra J. Kinney assisted in the preparation of the document and the coordination of the reviewers' comments.